

# GROUP ACTION AND $L^2$ -NORM ESTIMATES OF GEOMETRIC PROBLEMS

THANG PHAM

ABSTRACT. In 2017, by using the group theoretic approach, Bennett, Hart, Iosevich, Pakianathan, and Rudnev obtained a number of results on the distribution of simplices and sum-product type problems. The main purpose of this paper is to give a series of new applications of their powerful framework, namely, we focus on the product and quotient of distance sets, the  $L^2$ -norm of the direction set, and the  $L^2$ -norm of scales in difference sets.

## 1. INTRODUCTION

Let  $\mathbb{F}_q$  be a finite field of order  $q$ , where  $q$  is a prime power. Let  $O(d)$  be the orthogonal group of  $d$  by  $d$  matrices with entries in  $\mathbb{F}_q$ .

Given an integer  $k \geq 1$ , we say that two  $k$ -simplices in  $\mathbb{F}_q^d$  with vertices  $(x_1, \dots, x_{k+1})$  and  $(y_1, \dots, y_{k+1})$  are in the same congruence class if  $\|x_i - x_j\| = \|y_i - y_j\|$  for all  $1 \leq i \neq j \leq k+1$ . Here  $\|x_i - x_j\| = (x_{i1} - x_{j1})^2 + \dots + (x_{id} - x_{jd})^2$ . This is equivalent to say that there exist  $\theta \in O(d)$  and  $z \in \mathbb{F}_q^d$  such that  $y_i = \theta x_i + z$  for all  $1 \leq i \leq k+1$ . Given a set  $E \subset \mathbb{F}_q^d$ , we denote the set of congruence classes of  $k$ -simplices determined by  $E$  by  $T_k^d(E)$ .

The question of finding the smallest threshold  $\alpha$  such that  $|T_k^d(E)| \gg q^{\binom{k+1}{2}}$  whenever  $|E| \gg q^\alpha$  has a rich history, for instance, see [9, 24, 10, 3, 21]. The best current result is due to Bennett, Hart, Iosevich, Pakianathan, and Rudnev in [2]. More precisely, by using the group theoretic approach, they proved that for all  $1 \leq k \leq d$ , if  $|E| \gg q^{d - \frac{d-1}{k+1}}$ , then  $|T_k^d(E)| \gg q^{\binom{k+1}{2}}$ . In two dimensions, they are able to obtain better exponents, namely,  $8/5$  for  $k = 2$  and  $4/3$  for  $k = 1$ . We also note that when  $k = 1$ , this problem is known as the Erdős-Falconer distance problem in the literature, we refer the interested reader to [20, 15] for the recent progress.

It is worth noting that the group action techniques are not only useful in discrete setting, but it is also very powerful in the continuous setting. One example, we have to mention here, is an  $L^2$  identity due to Liu [17], which plays an important role in the recent breakthrough on the pinned Falconer distance problem [5, 6]. Similar results of this approach can also be found in [7, 8, 18].

The main purpose of this paper is to provide a series of new applications of this group theoretic approach, namely, on the product and quotient of distance sets, the  $L^2$ -norm of the direction set, and the  $L^2$ -norm of scales in difference sets.

Throughout this paper, we denote the set of square elements in  $\mathbb{F}_q$  by  $(\mathbb{F}_q)^2$ , and by  $X \gg Y$  we mean there exists an absolutely positive constant  $C$  such that  $X \geq CY$ .

**1.1. Product and Quotient of distance sets.** For  $E \subset \mathbb{F}_q^2$ , the distance set  $\Delta(E)$  is the set of all distances determined by pairs of points in  $E$ , namely,

$$\Delta(E) := \{|x - y| : x, y \in E\}.$$

We define the product, quotient, and sum of  $\Delta(E)$  as follows:

$$\frac{\Delta(E)}{\Delta(E)} := \left\{ \frac{a}{b} : a, b \in \Delta(E) \right\}, \quad \Delta(E) \cdot \Delta(E) := \{a \cdot b : a, b \in \Delta(E)\},$$

and

$$\Delta(E) + \Delta(E) := \{a + b : a, b \in \Delta(E)\}.$$

Our first result is on the size of  $\Delta(E) \cdot \Delta(E)$ .

**Theorem 1.1.** *For  $E \subset \mathbb{F}_q^2$  with  $|E| \gg q^{8/7}$ , we have*

$$|\Delta(E) \cdot \Delta(E)| \gg q.$$

For comparison, one can use graph theoretic techniques or discrete Fourier analysis to prove the same exponent for the sum set  $\Delta(E) + \Delta(E)$ , for instance, see [22]. It is necessary to say that the study of the product set would be much harder than the case of the sum set, the main reason comes from the observation that  $\Delta(E) + \Delta(E) = \Delta(E \times E)$ , so the sum set can be reduced to the original distance problem for Cartesian product sets, which possesses some nicely additive structures. As a consequence, when  $q$  is a prime, it has been shown in [4] that the exponent  $8/7$  can be improved to  $11/10$ . For the sum of two different distance sets, we refer the reader to [14] for more details. We also want to emphasize here that it is difficult to extend the methods from [4, 14, 22] to study the case of product set  $\Delta(E) \cdot \Delta(E)$ . As in the Erdős-Falconer distance problem, the conjectured exponent for product of distance sets should be 1.

For quotient of distance sets, by using discrete Fourier analysis, Iosevich, Koh, and Parshall [11] proved that if  $|E| \geq 9q$ , then

$$(1.1) \quad \frac{\Delta(E)}{\Delta(E)} = \mathbb{F}_q.$$

They also constructed examples to show that the condition  $|E| \gg q$  is optimal. There are two perspectives we want to mention here: their proof is very sophisticated, and it only tells us that the quotient contains the whole field, so given an element  $r \in \mathbb{F}_q$ , it is

not clear how many tuples  $(a, b, c, d) \in E^4$  such that  $\|a - b\|/\|c - d\| = r$ . Using the group theoretic method, we prove the following.

**Theorem 1.2.** *For  $E \subset \mathbb{F}_q^2$  with  $q \equiv 3 \pmod{4}$ . Assume that  $|E| \gg q$ , then for each non-zero square  $r$  in  $\mathbb{F}_q$ , the number of quadruples  $(a, b, c, d) \in E^4$  such that  $\|a - b\|/\|c - d\| = r$  is at least  $\gg |E|^4 q^{-1}$ . In particular,*

$$(\mathbb{F}_q)^2 \subseteq \frac{\Delta(E)}{\Delta(E)}.$$

We remark that the condition  $q \equiv 3 \pmod{4}$  is needed in the proof of this theorem, which helps to avoid pairs of zero distances.

**1.2. Distribution of directions.** Given  $E \subset \mathbb{F}_q^2$ , the  $L^2$ -norm of the direction set bounds the number of quadruples  $(u, v, x, y) \in E^4$  such that

$$(1.2) \quad (u - v) = \lambda \cdot (x - y), \quad x \neq y, \quad u \neq v,$$

for some  $\lambda \neq 0$ . We denote the number of such quadruples by  $L^2(D_E)$ . In the next theorem, we give an upper bound for this quantity.

**Theorem 1.3.** *For  $E \subset \mathbb{F}_q^2$ , we have*

$$\left| L^2(D_E) - \frac{|E|^4}{q} \right| \ll q^2 |E|^2.$$

If we are interested in the number of directions spanned by a set  $E \subset \mathbb{F}_q^2$ , then Theorem 1.3 says that we have at least  $\gg q$  directions as long as  $|E| \gg q^{3/2}$ . However, it is known in the literature that the condition  $|E| \gg q$  would be sufficient, see [12] for example. When the size of  $E$  is very small, say  $|E| \ll q$ , we refer the reader to [23] for related results.

**1.3. Scales in difference sets.** For  $E \subset \mathbb{F}_q^2$ , an element  $\lambda \in \mathbb{F}_q$  is called a scale in the difference set  $E - E$  if there exist  $u_1, v_1, u_2, v_2 \in E$  such that

$$(u_1 - v_1) = \lambda(u_2 - v_2).$$

We denote the set of all scales in  $E - E$  by  $\frac{E-E}{E-E}$ . In this paper, we are interested in bounding the  $L^2$ -norm, i.e. the number of tuples  $(u_1, v_1, u_2, v_2, u_3, v_3, u_4, v_4) \in E^8$  such that

$$(u_1 - v_1) = \lambda(u_2 - v_2), \quad (u_3 - v_3) = \lambda(u_4 - v_4),$$

for some  $\lambda \neq 0$ . We denote the number of such tuples by  $L^2(S_E)$ .

With the same approach, we have an upper bound for the  $L^2$ -norm of scales.

**Theorem 1.4.** *For  $E \subset \mathbb{F}_q^2$ , we have*

$$\left| L^2(S_E) - \frac{|E|^8}{q^3} \right| \ll |E|^6 + q^2 |E|^5.$$

For the set of scales, it is not hard to show that if  $|E| \gg q^{3/2}$ , then  $E - E/E - E$  covers the whole field. Roughly speaking, under the condition  $|E| \gg q^{3/2}$ , one can find a line with at least  $q^{1/2}$  points from  $E$ , then without loss of generality, we assume that line is defined by  $y = 0$ . So the problem is reduced to one dimensional version, namely,  $A - A/A - A$  for some  $A \subset \mathbb{F}_q$ . It is well-known that  $A - A/A - A = \mathbb{F}_q$  when  $|A| \gg q^{1/2}$ , see [1, Lemma 1] for example. We leave this to the reader for checking in detail. The exponent  $3/2$  is also optimal, for example, take  $q = p^2$  and  $E = \mathbb{F}_p \times \mathbb{F}_q$ , then  $|\frac{E-E}{E-E}| = p = o(q)$ .

It follows from Theorems 1.3 and 1.4 that  $L^2(D_E) = (1 + o(1))|E|^4/q$  and  $L^2(S_E) = (1 + o(1))|E|^8/q^3$  whenever  $|E| \gg q^{3/2}$  and  $|E| \gg q^{5/3}$ , respectively. We do not believe these two theorems are sharp, and offer the following conjectures.

**Conjecture 1.5.** *Let  $E$  be a subset of  $\mathbb{F}_q^2$ . If  $|E| \gg q^{1+\epsilon}$  for any  $\epsilon > 0$ , then*

$$L^2(D_E) = (1 + o(1))\frac{|E|^4}{q}.$$

**Conjecture 1.6.** *Let  $E$  be a subset of  $\mathbb{F}_q^2$ . If  $|E| \gg q^{\frac{3}{2}+\epsilon}$  for any  $\epsilon > 0$ , then*

$$L^2(S_E) = (1 + o(1))\frac{|E|^8}{q^3}.$$

The final remark is that all results in this paper can be extended to higher dimensions in the same way, but the proofs will become much complicated, for instance, in the proofs of Theorems 1.1 and 1.2, one has to count the number of pairs of zero distances, which can be done by using a number of results from Restriction theory [13]. Therefore, to keep this paper simple, we only present results in two dimensions.

## 2. NOTATIONS FROM DISCRETE FOURIER ANALYSIS

In this section, we recall some notations and results from discrete Fourier analysis which will be needed for our coming proofs in next sections.

Given a complex function  $f: \mathbb{F}_q^d \rightarrow \mathbb{C}$ , the Fourier transform of  $f$  is defined by

$$\widehat{f}(\xi) = \frac{1}{q^d} \sum_{x \in \mathbb{F}_q^d} f(x) \chi(-x \cdot \xi),$$

where  $\chi$  is a fixed non-trivial additive character of  $\mathbb{F}_q$ .

Using the orthogonality property of  $\chi$ , i.e.

$$\sum_{x \in \mathbb{F}_q^d} \chi(x \cdot \xi) = \begin{cases} 0 & \text{if } \xi \neq 0, \\ q^d & \text{if } \xi = 0, \end{cases}$$

one can prove that

$$f(x) = \sum_{\xi \in \mathbb{F}_q^d} \widehat{f}(\xi) \chi(\xi \cdot x).$$

In this setting, the Plancherel formula reads as

$$\sum_{\xi \in \mathbb{F}_q^d} |\widehat{f}(\xi)|^2 = q^{-d} \sum_{x \in \mathbb{F}_q^d} |f(x)|^2.$$

When  $f$  is the indicator of a given set  $E \subset \mathbb{F}_q^d$ , one has

$$\sum_{\xi \in \mathbb{F}_q^d} |\widehat{E}(\xi)|^2 = q^{-d} |E|.$$

The following lemma is known as the finite field analog of the spherical average in the classical Falconer distance problem [19, Chapter 3].

**Lemma 2.1.**

$$\max_{t \in \mathbb{F}_q \setminus \{0\}} \sum_{\|\eta\|=t} |\widehat{E}(\eta)|^2 \ll \frac{|E|^{3/2}}{q^3}.$$

A proof of this lemma can be found in [3, Lemma 4.4].

### 3. PROOF OF THEOREM 1.1

It is well-known that for  $u, v, x, y \in \mathbb{F}_q^d$ , if  $\|u - v\| = \|x - y\|$ , then there exist  $\theta \in O(d)$  and  $z \in \mathbb{F}_q^d$  such that

$$\theta u + z = x, \quad \theta v + z = y.$$

We now show that a similar statement holds for the case of product of distance sets.

**Lemma 3.1.** *Set*

$$G_1 = \left\{ \text{diag}(r_1 \theta_1, r_2 \theta_2) \in M_{2d \times 2d} : r_1 \cdot r_2 = 1, \theta_1, \theta_2 \in O(d) \right\},$$

and  $G = G_1 \times \mathbb{F}_q^{2d}$ . For  $x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4 \in \mathbb{F}_q^d$ , if

$$\|x_1 - y_1\| \cdot \|x_2 - y_2\| = \|x_3 - y_3\| \cdot \|x_4 - y_4\| \neq 0, \quad \frac{\|x_4 - y_4\|}{\|x_2 - y_2\|} \in (\mathbb{F}_q)^2,$$

then there exists  $g \in G$  such that

$$g(x_1, x_2)^T = (x_3, x_4)^T, \quad g(y_1, y_2)^T = (y_3, y_4)^T.$$

*Proof.* If

$$\|x_1 - y_1\| \cdot \|x_2 - y_2\| = \|x_3 - y_3\| \cdot \|x_4 - y_4\| \neq 0,$$

then

$$\frac{\|x_1 - y_1\|}{\|x_3 - y_3\|} = \frac{\|x_4 - y_4\|}{\|x_2 - y_2\|}.$$

Set  $r = \frac{\|x_4 - y_4\|}{\|x_2 - y_2\|}$ . From our assumptions, we know that  $r$  is a square, say, say,  $r = t^2$  for some  $t \neq 0$ . This infers that

$$\|(1/t)x_1 - (1/t)y_1\| = \|x_3 - y_3\|.$$

Hence, we can find  $\theta_1 \in O(d)$  and  $z_1 \in \mathbb{F}_q^d$  such that

$$(1/t)\theta_1 x_1 + z_1 = x_3, \quad (1/t)\theta_1 y_1 + z_1 = y_3.$$

Similarly, from the fact that

$$\|x_4 - y_4\| = \|tx_2 - ty_2\|,$$

we can find  $\theta_2 \in O(d)$  and  $z_2 \in \mathbb{F}_q^d$  such that

$$t\theta_2 x_2 + z_2 = x_4, \quad t\theta_2 y_2 + z_2 = y_4.$$

In other words,  $g = (1/t\theta_1, t\theta_2) \times (z_1, z_2) \in G$  is the element we want to find.  $\square$

*Proof of Theorem 1.1.* Let

$$A = \{(a, b) \in E \times E : \|a - b\| \in (\mathbb{F}_q)^2 \setminus \{0\}\},$$

and

$$B = E \times E \setminus (A \cup \{(a, b) \in E \times E : \|a - b\| = 0\}).$$

Let  $N_0(E)$  be the number of pairs  $(a, b) \in E \times E$  such that  $\|a - b\| = 0$ . We know from [16, Proposition 2.4] that

$$N_0(E) \ll \frac{|E|^2}{q} + q|E| \ll |E|^2,$$

when  $|E| \gg q$ . Thus, either  $|A|$  or  $|B|$  is bounded from below by  $|E|^2/2$ . Without loss of generality, we assume that  $|B| \gg |E|^2/2$ . For any  $\lambda \in (\mathbb{F}_q)^2 \setminus \{0\}$ , define

$$\nu(\lambda) := \#\{(x_1, y_1, x_2, y_2) \in B^2 : \|x_1 - y_1\| \cdot \|x_2 - y_2\| = \lambda\}.$$

We have

$$\sum_{\lambda \in (\Delta(E) \cdot \Delta(E)) \cap (\mathbb{F}_q)^2 \setminus \{0\}} \nu(\lambda) \gg |E|^4.$$

By the Cauchy-Schwarz inequality, one has

$$\sum_{\lambda \in \Delta(E) \cap (\mathbb{F}_q)^2 \setminus \{0\}} \nu(\lambda) \ll |\Delta(E)|^{1/2} \cdot \left( \sum_{\lambda \in \mathbb{F}_q} \nu(\lambda)^2 \right)^{1/2}.$$

In the next step, we are going to bound  $\sum_{\lambda \in \mathbb{F}_q} \nu(\lambda)^2$  from above. For  $\theta = \text{diag}(r_1\theta_1, r_2\theta_2) \in G_1$  and  $z = (z_1, z_2) \in \mathbb{F}_q^2 \times \mathbb{F}_q^2$ , we define  $\mu_\theta(z)$  to be the number of tuples  $(x_1, x_2, x_3, x_4) \in E^4$  such that

$$\theta(x_1, x_2)^T + (z_1, z_2)^T = (x_3, x_4)^T.$$

Then we observe that

$$\sum_{\lambda} \nu(\lambda)^2 \leq \sum_{\theta \in G_1, z \in \mathbb{F}_q^4} \mu_\theta(z)^2.$$

It suffices to prove that

$$\sum_{\theta \in G_1, z \in \mathbb{F}_q^4} \mu_\theta(z)^2 \ll \frac{|E|^8}{q},$$

whenever  $|E| \gg q^{8/7}$ .

Set  $F = E \times E$ . One has

$$\begin{aligned} \mu_\theta(z) &= \sum_{x_1, x_2 \in \mathbb{F}_q^2} F(x_1, x_2) F(\theta(x_1, x_2)^T + z) \\ &= \sum_{x_1, x_2 \in \mathbb{F}_q^2} F(x_1, x_2) \sum_{m \in \mathbb{F}_q^4} \widehat{F}(m) \chi(m \cdot (\theta(x_1, x_2)^T + z)) \\ &= q^4 \sum_{m \in \mathbb{F}_q^4} \widehat{F}(m) \widehat{F}(-\theta^T m) \chi(m \cdot z). \end{aligned}$$

Hence,

$$\widehat{\mu}_\theta(\xi) = q^4 \widehat{F}(-\xi) \widehat{F}(\theta^T \xi).$$

Therefore, for a fixed  $\theta \in G_1$ ,

$$\sum_{\xi \in \mathbb{F}_q^4} |\widehat{\mu}_\theta(\xi)|^2 = \frac{|F|^4}{q^8} + \sum_{\xi \neq 0} |\widehat{\mu}_\theta(\xi)|^2.$$

By writing  $\xi = (\xi_1, \xi_2) \in \mathbb{F}_q^2 \times \mathbb{F}_q^2$  and recall that  $F = E \times E$ , we have

$$\begin{aligned} \sum_{\theta \in G_1, \xi \in \mathbb{F}_q^4 \setminus \{0\}} |\widehat{\mu}_\theta(\xi)|^2 &= q^8 \sum_{\theta, \xi \neq 0} |\widehat{F}(-\xi)|^2 |\widehat{F}(\theta^T \xi)|^2 \\ &= q^8 \sum_{\theta} \sum_{\xi_1=0, \xi_2 \neq 0} + q^8 \sum_{\theta} \sum_{\xi \neq 0, \xi_2=0} + q^8 \sum_{\theta} \sum_{\xi_1 \neq 0, \xi_2 \neq 0} =: I + II + III. \end{aligned}$$

For  $I$ , by Plancherel, one has

$$\begin{aligned} I &= q^8 \sum_{\theta_1, \theta_2 \in O(2)} \sum_{r_1, r_2: r_1 r_2 = 1} \sum_{\xi_1=0, \xi_2 \neq 0} |\widehat{E}(-\xi_1)|^2 |\widehat{E}(-\xi_2)|^2 |\widehat{E}(r_1 \theta_1^T \xi_1)|^2 |\widehat{E}(r_2 \theta_2^T \xi_2)|^2 \\ &= q |E|^4 \sum_{\theta_2} \sum_{r_2 \neq 0} \sum_{\xi_2 \neq 0} |\widehat{E}(-\xi_2)|^2 |\widehat{E}(r_2 \theta_2^T \xi_2)|^2 \\ &= q |E|^4 \sum_{\xi_2 \neq 0} |\widehat{E}(-\xi_2)|^2 \sum_{r \neq 0} \sum_{\|\eta\| = \|\xi_2\|} |\widehat{E}(r\eta)|^2 \\ &\ll q^{-3} |E|^6. \end{aligned}$$

Similarly,  $II \ll q^{-3} |E|^6$ . For  $III$ , we proceed as follows:

$$\begin{aligned} III &= q^8 \sum_{\theta} \sum_{r_1, r_2: r_1 r_2 = 1} \sum_{\xi_1 \neq 0, \xi_2 \neq 0} |\widehat{E}(-\xi_1)|^2 |\widehat{E}(-\xi_2)|^2 |\widehat{E}(r_1 \theta_1^T \xi_1)|^2 |\widehat{E}(r_2 \theta_2^T \xi_2)|^2 \\ &= q^8 \sum_{\theta_2} \sum_{\xi_2 \neq 0} \sum_{r_2 \neq 0} |\widehat{E}(\xi_2)|^2 |\widehat{E}(r_2 \theta_2^T \xi_2)|^2 \sum_{\xi_1 \neq 0} |\widehat{E}(\xi_1)|^2 \sum_{\eta: \|\eta\| = \frac{\|\xi_1\|}{r_2^2}} |\widehat{E}(\eta)|^2. \end{aligned}$$

This implies

$$III \ll q^8 \left( \sum_{\xi_1} |\widehat{E}(\xi_1)|^2 \right)^3 \cdot \left( \max_{t \in \mathbb{F}_q \setminus \{0\}} \sum_{\|\eta\|=t} |\widehat{E}(\eta)|^2 \right).$$

Using Lemma 2.1, we obtain

$$III \ll q^{-1} |E|^{9/2}.$$

In other words,

$$\sum_{\theta, \xi} |\widehat{\mu}_\theta(\xi)|^2 \ll \frac{|E|^8}{q^5} + \frac{|E|^6}{q^3} + \frac{|E|^{9/2}}{q}.$$

This infers

$$\sum_{\theta \in G_1, z \in \mathbb{F}_q^4} \mu_\theta(z)^2 \ll \frac{|E|^8}{q} + q|E|^6 + q^3|E|^{9/2} \ll \frac{|E|^8}{q},$$

whenever  $|E| \gg q^{8/7}$ . □

#### 4. PROOF OF THEOREM 1.2

We note that  $r = 0$  is trivially in  $\Delta(E)/\Delta(E)$ . For  $\theta \in O(2)$  and  $z \in \mathbb{F}_q^2$ , assume  $r$  is a nonzero square, we define

$$\eta_\theta(z) := \#\{(u, v) \in E \times E : u - \sqrt{r}\theta v = z\}.$$

We first observe that

$$\sum_{\theta \in O(2), z \in \mathbb{F}_q^2} \eta_\theta(z) = |E|^2 \cdot |O(2)|.$$

So by the Hölder inequality, one has

$$\sum_{\theta, z} \eta_\theta(z)^2 \geq \frac{|E|^4 |O(2)|}{q^2}.$$

On the other hand, the sum  $\sum_{\theta, z} \eta_\theta(z)^2$  counts the number of tuples  $(a, b, c, d) \in E^4$  such that  $a - \sqrt{r}\theta c = z$  and  $b - \sqrt{r}\theta d = z$ , so  $\|a - b\| = r\|c - d\|$ . For such a tuple, we have  $r \in \Delta(E)/\Delta(E)$ , unless  $\|a - b\| = \|c - d\| = 0$ .

For each tuple  $(a, b, c, d) \in E^4$  such that  $\|a - b\| = r\|c - d\|$ , we have two possibilities.

If  $a - b \neq 0$ , then this tuple is counted  $\sim |O(1)| \sim 1$  times in the sum  $\sum_{\theta, z} \eta_\theta(z)^2$ , since the size of the stabilizer of each non-zero element is  $\sim |O(1)|$ .

If  $a = b$ , then  $c = d$ . Moreover, the number of tuples  $(a, b, c, d) \in E^4$  with  $a = b$  and  $c = d$  in the sum  $\sum_{\theta, z} \eta_\theta(z)^2$  is equal to  $\sum_{\theta, z} \eta_\theta(z) = |E|^2 |O(2)|$ .

From these observations, the number of tuples  $(a, b, c, d) \in E^4$  such that  $\|a - b\| = r\|c - d\| \neq 0$  is at least

$$\frac{1}{|O(1)|} \left( \sum_{\theta, z} \eta_\theta(z)^2 - |E|^2 |O(2)| - |O(1)| N_0 \right),$$

where  $N_0$  is the number of tuples with  $\|a - b\| = r\|c - d\| = 0$ ,  $a \neq b$ , and  $c \neq d$ .



Since  $q \equiv 3 \pmod{4}$ , it is clear that  $N_0 = 0$ . Thus,

$$\frac{1}{|O(1)|} \left( \sum_{\theta, z} \eta_{\theta}(z)^2 - |E|^2 |O(2)| - |O(1)| N_0 \right) \gg \frac{|E|^4}{q},$$

whenever  $|E| \gg q$ .

### 5. PROOF OF THEOREM 1.3

Define the group  $G = \mathbb{F}_q \times \mathbb{F}_q^2$  with the operator

$$(\lambda, z_1) * (\beta, z_2) = (\lambda \cdot \beta, z_1 + z_2).$$

For a fixed  $\lambda$ , we define the counting function

$$\gamma_{\lambda}(z) := \#\{(a, b) \in E^2 : \lambda a + z = b\}.$$

By a direct computation, the tuple  $(u, v, x, y)$  satisfies (1.2) if and only if there exist  $\lambda \neq 0$  and  $z \in \mathbb{F}_q^2$  such that  $\lambda u + z = x$  and  $\lambda v + z = y$ . Thus,

$$L^2(S_E) = \sum_{\lambda, z} \gamma_{\lambda}(z)^2.$$

For a fixed  $\lambda$ , we have

$$\begin{aligned} \sum_{z \in \mathbb{F}_q^2} \gamma_{\lambda}(z)^2 &= q^2 \sum_{\xi \in \mathbb{F}_q^2} |\widehat{\gamma_{\lambda}}(\xi)|^2 = \frac{|\sum_z \gamma_{\lambda}(z)|^2}{q^2} + q^2 \sum_{\xi \neq 0} |\widehat{\gamma_{\lambda}}(\xi)|^2 \\ &= \frac{|E|^4}{q^2} + q^2 \sum_{\xi \neq 0} |\widehat{\gamma_{\lambda}}(\xi)|^2. \end{aligned}$$

On the other hand,

$$\begin{aligned} \gamma_{\lambda}(z) &= \sum_u E(u) E(\lambda u + z) = \sum_u E(u) \sum_{\xi} \widehat{E}(\xi) \chi(\xi \cdot (\lambda u + z)) \\ &= \sum_{\xi} \widehat{E}(\xi) \chi(\xi \cdot z) \sum_u E(u) \chi(\lambda \xi \cdot u) \\ &= q^2 \sum_{\xi} \widehat{E}(\xi) \widehat{E}(-\lambda \xi) \chi(\xi \cdot z). \end{aligned}$$

Hence,

$$\widehat{\gamma_{\lambda}}(\xi) = q^2 \widehat{E}(-\xi) \widehat{E}(\lambda \xi).$$

So,

$$\sum_{\xi \neq 0} |\widehat{\gamma_{\lambda}}(\xi)|^2 = q^4 \sum_{\xi \neq 0} |\widehat{E}(-\xi)|^2 |\widehat{E}(\lambda \xi)|^2.$$

Hence,

$$\sum_{\lambda \neq 0} \sum_{\xi \neq 0} |\widehat{\gamma_{\lambda}}(\xi)|^2 = q^4 \sum_{\xi \neq 0} |\widehat{E}(\xi)|^2 \sum_{m \in \langle \xi \rangle \setminus \{0\}} |\widehat{E}(m)|^2,$$

where  $\langle \xi \rangle$  is the line generated by  $\xi$ .

To proceed further, we set  $L := \langle \xi \rangle$  and  $T_L(\mathbf{E}) := \sum_{m \in L} |\widehat{\mathbf{E}}(m)|^2$ . It follows that

$$T_L(\mathbf{E}) = q^{-4} \sum_{m \in L} \sum_{x, y \in \mathbf{E}} \chi(m \cdot (x - y)) = q^{-4} \sum_{x, y \in \mathbf{E}} \sum_{t \in \mathbb{F}_q} \chi(t(\xi \cdot (x - y))) = q^{-4}(I + II),$$

where

$$I = \sum_{x, y \in \mathbf{E}, \xi \cdot (x - y) = 0} \sum_{t \in \mathbb{F}_q} \chi(t(\xi \cdot (x - y))) = q \sum_{x, y \in \mathbf{E}, \xi \cdot (x - y) = 0} 1,$$

and

$$II = \sum_{x, y \in \mathbf{E}, \xi \cdot (x - y) \neq 0} \sum_{t \in \mathbb{F}_q} \chi(t(\xi \cdot (x - y))) = 0.$$

We note that the equation  $\xi \cdot (x - y) = 0$  with  $x, y \in \mathbf{E}$  means that  $x - y$  lies on the line with the normal vector  $\xi$  and passing through the origin. Using the fact that each line in  $\mathbb{F}_q^2$  contains exactly  $q$  points, we have  $I \leq q^2 |\mathbf{E}|$ . Thus, using Plancherel formula, we obtain

$$\sum_{\lambda \neq 0} \sum_{\xi \neq 0} |\widehat{\gamma_\lambda(\xi)}|^2 \leq |\mathbf{E}|^2.$$

In other words,

$$\left| \sum_{\lambda, z} \gamma_\lambda(z)^2 - \frac{|\mathbf{E}|^4}{q} \right| \ll q^2 |\mathbf{E}|^2.$$

This completes the proof of the theorem.

## 6. PROOF OF THEOREM 1.4

Using the notations as in the previous section, we observe that

$$L^2(S_{\mathbf{E}}) = \sum_{\lambda \neq 0} \left( \sum_z \gamma_\lambda(z)^2 \right)^2.$$

As in the previous section, we know that

$$\sum_{z \in \mathbb{F}_q^2} \gamma_\lambda(z)^2 = \frac{|\mathbf{E}|^4}{q^2} + q^2 \sum_{\xi \neq 0} |\widehat{\gamma_\lambda(\xi)}|^2,$$

and

$$\sum_{\lambda} \sum_{\xi \neq 0} |\widehat{\gamma_\lambda(\xi)}|^2 \ll |\mathbf{E}|^2.$$

Thus,

$$\left| \sum_{\lambda \neq 0} \left( \sum_z \gamma_\lambda(z)^2 \right)^2 - \frac{|\mathbf{E}|^8}{q^3} \right| \ll |\mathbf{E}|^6 + q^4 \sum_{\lambda} \left( \sum_{\xi \neq 0} |\widehat{\gamma_\lambda(\xi)}|^2 \right)^2.$$

As we computed in the previous section, one has

$$\left( \sum_{\xi \neq 0} |\widehat{\gamma_\lambda(\xi)}|^2 \right)^2 = q^8 \left( \sum_{\xi \neq 0} |\widehat{\mathbf{E}}(\xi)|^2 |\widehat{\mathbf{E}}(-\lambda\xi)|^2 \right)^2.$$

Therefore,

$$\begin{aligned} \sum_{\lambda} \left( \sum_{\xi \neq 0} |\widehat{E}(\xi)|^2 |\widehat{E}(-\lambda\xi)|^2 \right)^2 &= \sum_{\xi_1 \neq 0, \xi_2 \neq 0} |\widehat{E}(\xi_1)|^2 |\widehat{E}(\xi_2)|^2 \sum_{\lambda} |\widehat{E}(\lambda\xi_1)|^2 |\widehat{E}(\lambda\xi_2)|^2 \\ &= \sum_{\xi_1 \neq 0, \xi_2 \neq 0} |\widehat{E}(\xi_1)|^2 |\widehat{E}(\xi_2)|^2 \cdot L(\xi_1, \xi_2), \end{aligned}$$

where

$$L(\xi_1, \xi_2) = \sum_{\lambda} |\widehat{E}(\lambda\xi_1)|^2 |\widehat{E}(\lambda\xi_2)|^2.$$

To proceed further, we need the following lemma.

**Lemma 6.1.** *For  $\xi_1, \xi_2 \neq 0$ , one has*

$$L(\xi_1, \xi_2) = \sum_{\lambda} |\widehat{E}(\lambda\xi_1)|^2 |\widehat{E}(\lambda\xi_2)|^2 \ll \frac{|E|^3}{q^6}.$$

*Proof.* By the Cauchy-Schwarz inequality, we have

$$L(\xi_1, \xi_2) \leq \left( \sum_{\lambda} |\widehat{E}(\lambda\xi_1)|^4 \right)^{1/2} \cdot \left( \sum_{\lambda} |\widehat{E}(\lambda\xi_2)|^4 \right)^{1/2}.$$

Moreover, for  $m \neq 0$ ,

$$\begin{aligned} \sum_{\lambda} |\widehat{E}(\lambda m)|^4 &= \frac{1}{q^8} \sum_{x, y, z, w \in E} \sum_{\lambda} \chi(m\lambda(x+y-z-w)) \\ &= \frac{1}{q^7} \sum_{x, y, z, w \in E} \mathbf{1}_{m \cdot (x+y-z-w)=0} \\ &\ll \frac{|E|^3}{q^6}, \end{aligned}$$

where the last inequality follows from the number of tuples  $(x, y, z, w) \in E^4$  such that  $x+y-z-w$  lies on the line with the normal vector  $m$  and passing through the origin.

This completes the proof.  $\square$

To continue, we apply the Plancherel formula,

$$\begin{aligned} \sum_{\lambda} \left( \sum_{\xi \neq 0} |\widehat{E}(\xi)|^2 |\widehat{E}(-\lambda\xi)|^2 \right)^2 &= \sum_{\xi_1 \neq 0, \xi_2 \neq 0} |\widehat{E}(\xi_1)|^2 |\widehat{E}(\xi_2)|^2 \cdot L(\xi_1, \xi_2) \\ &\ll \frac{|E|^5}{q^{10}}. \end{aligned}$$

In other words, we obtain

$$\left| \sum_{\lambda} \left( \sum_z \gamma_{\lambda}(z)^2 \right)^2 - \frac{|E|^8}{q^3} \right| \ll |E|^6 + q^2 |E|^5.$$

## 7. ACKNOWLEDGEMENT

The author would like to thank to Vietnam Institute for Advanced Study in Mathematics (VIASM) for the hospitality and for the excellent working condition.

## REFERENCES

- [1] A. Balog, *Another sum-product estimate in finite fields*, Proceedings of the Steklov Institute of Mathematics, **280**(2) (2013): 23–29.
- [2] M. Bennett, D. Hart, A. Iosevich, J. Pakianathan, and M. Rudnev, *Group actions and geometric combinatorics in  $\mathbb{F}_q^d$* , Forum Mathematicum, **29**(1) (2017): 91–110.
- [3] J. Chapman, M. Burak Erdogan, D. Hart, A. Iosevich, and D. Koh, *Pinned distance sets,  $k$ -simplices, Wolff’s exponent in finite fields and sum-product estimates*, Mathematische Zeitschrift, **271** (1-2) (2012): 63–93.
- [4] D. Cheong, D. Koh, and T. Pham, *An asymmetric bound for sum of distance sets*, Proceedings of the Steklov Institute of Mathematics, **314**(1) (2021): 279–289.
- [5] X. Du, A. Iosevich, Y. Ou, H. Wang, and R. Zhang, *An improved result for Falconer’s distance set problem in even dimensions*, Mathematische Annalen, **380** (2021): 1215–1231.
- [6] L. Guth, A. Iosevich, Y. Ou, and H. Wang, *On Falconer’s distance set problem in the plane*, Inventiones mathematicae, **219**(3) (2022): 779–830.
- [7] A. Greenleaf, A. Iosevich, B. Liu, and E. Palsson, *A group-theoretic viewpoint on Erdős–Falconer problems and the Mattila integral*, Revista Matemática Iberoamericana, **31**(3) (2015): 799–810.
- [8] A. Greenleaf, A. Iosevich, and S. Mkrtychyan, *Existence of similar point configurations in thin subsets of  $\mathbb{R}^d$* , Mathematische Zeitschrift, **297**(1) (2021): 855–865.
- [9] D. Hart and A. Iosevich, *Ubiquity of simplices in subsets of vector spaces over finite fields*, Analysis Mathematica, **34** (2007).
- [10] D. Hart, A. Iosevich, D. Koh, S. Senger, and Ignacio Uriarte-Tuero, *Distance graphs in vector spaces over finite fields*, Recent advances in harmonic analysis and applications, pp. 139–160. Springer, New York, NY, 2012.
- [11] A. Iosevich, D. Koh, and H. Parshall, *On the quotient set of the distance set*, Moscow Journal of Combinatorics and Number Theory, **8**(2) (2019): 103–115.
- [12] A. Iosevich, H. Morgan, and J. Pakianathan, *On directions determined by subsets of vector spaces over finite fields*, Integers, 11:A39, 2011.
- [13] A. Iosevich, D. Koh, S. Lee, T. Pham, and C-Y. Shen, *On restriction estimates for the zero radius sphere over finite fields*, Canadian Journal of Mathematics, **73**(3)(2021): 769–786.
- [14] D. Koh, T. Pham, C-Y. Shen, and L. A. Vinh, *A sharp exponent on sum of distance sets over finite fields*, Mathematische Zeitschrift, **297**(3) (2021): 1749–1765.
- [15] D. Koh, T. Pham, and L. A. Vinh, *Extension theorems and a connection to the Erdős–Falconer distance problem over finite fields*, Journal of Functional Analysis, **281**(8) (2021): 109137.
- [16] D. Koh and H. Sun, *Distance sets of two subsets of vector spaces over finite fields*, Proceedings of the American Mathematical Society, **143**(4) (2015): 1679–1692.
- [17] B. Liu, *An  $L^2$ -identity and pinned distance problem*, Geometric and Functional Analysis, **29**(1) (2019): 283–294.
- [18] B. Liu, *Group actions, the Mattila integral and applications*, Proceedings of the American Mathematical Society, **147**(6) (2019): 2503–2516.
- [19] P. Mattila, *Fourier analysis and Hausdorff dimension*, Vol. 150, Cambridge University Press, 2015.
- [20] B. Murphy, G. Petridis, T. Pham, M. Rudnev, and S. Stevens, *On the pinned distances problem in positive characteristic*, Journal of the London Mathematical Society, **105**(1) (2022): 469–499.

- [21] H. Parshall, *Simplices over finite fields*, Proceedings of the American Mathematical Society, **145**(6) (2017): 2323–2334.
- [22] T. Pham, *Erdős distinct distances problem and extensions over finite spaces*, Phd thesis, EPFL, Lausanne Switzerland, 2017.
- [23] T. Szőnyi, *On the number of directions determined by a set of points in an affine Galois plane*, Journal of Combinatorial Theory Series A, **74**(1) (1996): 141–146.
- [24] L. A. Vinh, *On kaleidoscopic pseudo-randomness of finite Euclidean graphs*, Discussiones Mathematicae: Graph Theory, **32**(2) (2012).

HANOI UNIVERSITY OF SCIENCE, VIETNAM NATIONAL UNIVERSITY

*Email address:* thangpham.math@vnu.edu.vn