

On the existence and uniqueness of weak solutions to time-fractional elliptic equations with time dependent variable coefficients

H.T. Tuan*

November 30, 2020

Abstract

This paper is devoted to discuss on the existence and uniqueness of weak solutions to time-fractional elliptic equations having time dependent variable coefficients. To obtain the main result, our strategy is to combine Galerkin method, a basic inequality for fractional derivative of convex Lyapunov candidate functions, the Yoshida approximation sequence and the weak compactness argument.

Keywords and phases: Time fractional derivatives, Existence and uniqueness, Weak solution, Elliptic equations, Variable coefficients

1 Introduction

Diffusion equations with fractional-order derivatives in time (which is called as time-fractional diffusion equations) have been introduced in Physics by Nigmatullin [N86] to describe super slow diffusion process in a porous medium with the structure type of fractal geometry (Koch's tree). Then, by the probabilistic point of view, in the paper [MK2000], Metzler and Klafter have pointed out that a time-fractional diffusion equation generates a non-Markovian diffusion process with a long memory. After that, Roman and Alemany [RA94] have considered continuous-time random walks on fractals and observed that the average probability density of random walks on fractals obeys a diffusion equation with a fractional time derivative asymptotically. Another context where such systems appear is the modelling of evolution processes in materials with memory, see e.g., [P93, C99].

*httuan@math.ac.vn, Institute of Mathematics, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, 10307 Ha Noi, Viet Nam

The existence of solutions to time-fractional partial differential equations has been studied by many authors. In [EK04], using Fourier transform, the authors have obtained a fundamental solution for time-fractional elliptic equations with smooth coefficient. Combining Galerkin method and the Yoshida approximation sequence, in [Z09], the author has proposed a way to prove existence of certain weak solutions to abstract evolutionary integro-differential equations in Hilbert spaces. By virtue of the operator theory in functional analysis and the eigenfunction expansion method for symmetry elliptic operators, in [SY11], Sakamoto and Yamamoto have proved the existence and uniqueness of the weak solution for a fractional diffusion-wave equation. Based on a definition of the Caputo derivative on a finite interval in fractional Sobolev spaces, Gorenflo, Luchko and Yamamoto [GLY15] have investigated solutions (in the distribution sense) to time-fractional diffusion equations from the operator theoretic viewpoint. In the recent work, by a classical variational approach, K.V. Bockstal [B20] has established the existence of a unique weak solution to a class of fractional diffusion equations with Caputo derivative.

However, to the best of our knowledge, the development of this theory is still in its infancy and requires further researches. The main difficulty which one have to face is the meaning of the initial condition of solutions and the correctness of the formulation of weak solutions. In this paper, we focus on an initial-boundary value problem with the zero initial condition for time-fractional elliptic equations having time dependent variable coefficients which can not apply Fourier transformation to solve them. To overcome the aforementioned obstacles, our strategy is to use Galerkin method, a basic inequality for fractional derivative of convex Lyapunov candidate functions, the Yoshida approximation sequence and the weak compactness argument.

The paper is organized as follows. In section 2, we recall some preliminary results on fractional calculus. Then, we give the setting of the problem and propose a clear definition of a weak solution to a time fractional elliptic equation with time dependent variable coefficients. The main result of the paper is Theorem 8 on the existence and uniqueness of weak solutions introduced in section 3.

To conclude the introduction, we will introduce some notations used throughout the rest of the article. Denote by \mathbb{N} the set of natural numbers and by \mathbb{R} the set of real numbers. For any natural $d \in \mathbb{N}$, let \mathbb{R}^d be the d -dimensional Euclidean space. For a open subset Ω of \mathbb{R}^d , let $C_c^\infty(\Omega)$ denote the space of infinitely differentiable functions $f : \Omega \rightarrow \mathbb{R}$ with the compact support in Ω , $L^p(\Omega)$, $p \in \mathbb{N}$, be the set of measurable functions $f : \Omega \rightarrow \mathbb{R}$ such that

$$\int_{\Omega} |f(x)|^p dx < \infty,$$

$H^1(\Omega)$ be the Sobolev space containing all locally integrable functions $f : \Omega \rightarrow \mathbb{R}$ such that f and its weak derivatives belong to $L^2(\Omega)$, $H_0^1(\Omega)$ be the closure of $C_c^\infty(\Omega)$ in $H^1(\Omega)$, and $H^{-1}(\Omega)$ be the dual space of $H_0^1(\Omega)$. Fix $T > 0$, denote

by $W_1^1([0, T]; \mathbb{R})$ the space of functions $f : [0, T]; \mathbb{R}$ such that f and its weak derivative belong to the space $L^1([0, T]; \mathbb{R})$ and $W_2^1([0, T]; H^{-1}(\Omega))$ by the space of functions $f : [0, T] \rightarrow H^{-1}(\Omega)$ such that f and the weak derivative belong to the space $L^2([0, T]; H^{-1}(\Omega))$.

2 Fractional calculus

We briefly recall an abstract framework of fractional calculus.

Let $\alpha \in (0, 1]$, $[0, T] \subset \mathbb{R}$ and $x : [0, T] \rightarrow \mathbb{R}$ be a measurable function such that $\int_0^T |x(\tau)| d\tau < \infty$. The right-handed *Riemann–Liouville integral operator of order α* is defined by

$$(I_{0+}^\alpha x)(t) := \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} x(\tau) d\tau,$$

where $\Gamma(\cdot)$ is the Gamma function. The left-handed *Riemann–Liouville integral operator of order α* is defined by

$$(I_{T-}^\alpha x)(t) := \frac{1}{\Gamma(\alpha)} \int_t^T (\tau - t)^{\alpha-1} x(\tau) d\tau.$$

We have the following result on the relation between the right-handed and left-handed Riemann–Liouville integral operators.

Theorem 2.1 (The Hardy–Littlewood form of fractional integration by parts).
If $p > 1$, $q > 1$, $0 < \alpha < 1$, $\frac{1}{p} + \frac{1}{q} - 1 \leq \alpha$, and

$$f \in L^p([0, T]; \mathbb{R}), \quad g \in L^q([0, T]; \mathbb{R}),$$

then

$$\int_0^T (I_{0+}^\alpha f(t))g(t)dt = \int_0^T f(t)I_{T-}^\alpha g(t)dt.$$

Proof. See [LY37]. □

The right-handed *Riemann–Liouville fractional derivative* ${}^{RL}D_{\alpha+}^\alpha x$ of x on $[0, T]$ is defined by

$${}^{RL}D_{\alpha+}^\alpha x(t) := (DI_{0+}^{1-\alpha} x)(t) \text{ for almost } t \in [0, T],$$

where $D = \frac{d}{dt}$ is the usual derivative. The left-handed *Riemann–Liouville fractional derivative* ${}^{RL}D_{T-}^\alpha x$ of x on $[0, T]$ is defined by

$${}^{RL}D_{T-}^\alpha x(t) := -(DI_{T-}^{1-\alpha} x)(t) \text{ for almost } t \in [0, T].$$

Theorem 2.2. The formula

$$\int_0^T f(t)^{RL} D_{0+}^\alpha g(t) dt = \int_0^T g(t)^{RL} D_{T-}^\alpha f(t) dt$$

is valid for $0 < \alpha < 1$, $f \in I_{T-}^\alpha(L^p)$, $g \in I_{0+}^\alpha(L^q)$ and $\frac{1}{p} + \frac{1}{q} - 1 \leq \alpha$.

Proof. See [SKM93, Corollary 2, pp. 46]. □

The right-handed *Caputo fractional derivative* of x on $[0, T]$ is defined by

$${}^C D_{0+}^\alpha x(t) = {}^{RL} D_{a+}^\alpha (x(t) - x(0)) \quad \text{for almost } t \in [0, T]$$

and the left-handed *Caputo fractional derivative* of x on $[0, T]$ is defined by

$${}^C D_{T-}^\alpha x(t) = {}^{RL} D_{T-}^\alpha (x(t) - x(T)) \quad \text{for almost } t \in [0, T].$$

We have a sufficient condition for the existence of fractional derivative.

Theorem 2.3. Let $f \in AC([0, T]; \mathbb{R})$, $\alpha \in (0, 1)$, then ${}^{RL} D_{0+}^\alpha f$ and ${}^{RL} D_{T-}^\alpha f$ exist almost everywhere. Moreover, ${}^{RL} D_{0+}^\alpha f, {}^{RL} D_{T-}^\alpha f \in L^r([0, T]; \mathbb{R})$, $1 \leq r \leq \frac{1}{\alpha}$, and

$$\begin{aligned} {}^{RL} D_{0+}^\alpha f(t) &= \frac{1}{\Gamma(1-\alpha)} \left[\frac{f(0)}{t^\alpha} + \int_0^t \frac{f'(\tau)}{(t-\tau)^\alpha} d\tau \right], \\ {}^{RL} D_{T-}^\alpha f(t) &= \frac{1}{\Gamma(1-\alpha)} \left[\frac{f(T)}{(T-t)^\alpha} - \int_t^T \frac{f'(\tau)}{(\tau-t)^\alpha} d\tau \right]. \end{aligned}$$

Proof. See [SKM93, Lemma 2.2, pp. 35–36]. □

Definition 2.4. Let $u \in L^1([0, T]; H_0^1(\Omega))$. We define the weak Riemann–Liouville fractional derivative of the order α of u , ${}^{RL} D_{0+}^\alpha u(t)$, as below

$$\int_0^T \varphi(t) {}^{RL} D_{0+}^\alpha u(t) dt = \int_0^T {}^{RL} D_{T-}^\alpha \varphi(t) u(t) dt$$

for all $\varphi \in C_c^\infty((0, T); \mathbb{R})$.

3 Weak solutions to time-fractional elliptic equations

Let Ω be a bounded domain in \mathbb{R}^d with the boundary $\partial\Omega \in C^1$, $T > 0$ and $\alpha \in (0, 1)$. Denote $\Omega_T = (0, T] \times \Omega$. We consider the equation of the order α

$$\begin{aligned} \frac{\partial^\alpha u(t, x)}{\partial t^\alpha} - \sum_{i,j=1}^d \partial_{x_i} (a_{ij}(t, x) \partial_{x_j} u(t, x)) \\ + \sum_{j=1}^d b_j(t, x) \partial_{x_j} u(t, x) + c(t, x) u(t, x) = f(t) \end{aligned} \quad (1)$$

for $(t, x) \in \Omega_T$, where $\frac{\partial^\alpha u(t, \cdot)}{\partial t^\alpha}$ is the weak Riemann–Liouville fractional derivative of the order α of u with respect to the time variable t and

- (a1) $a_{ij}, b_j, c \in L^\infty(\Omega_T; \mathbb{R})$ for all $1 \leq i, j \leq d$;
- (a2) $a_{ij} = a_{ji}$ for all $1 \leq i, j \leq d$;
- (a3) there exists $\theta > 0$ such that $\sum_{i,j=1}^d a_{ij}(t, x) \xi_i \xi_j \geq \theta \|\xi\|^2$ for a.e. $t \in (0, T)$, $x \in \Omega$ and for all $\xi \in \mathbb{R}^d$;
- (a4) $f \in L^\infty([0, T]; H^{-1}(\Omega))$.

Assume that

$$u(t, x) = 0 \text{ on } [0, T] \times \partial\Omega. \quad (2)$$

Denote

$$a(u, v; t) := \int_{\Omega} - \sum_{1 \leq i, j \leq d} a_{ij} \partial_{x_j} u \partial_{x_i} v + \sum_{1 \leq j \leq d} b_j \partial_{x_j} uv + cuv$$

for all $u, v \in H_0^1(\Omega)$ and $\mathbf{u}(t) := [u(t)](x)$ for all $t \in [0, T]$, $x \in \Omega$.

Definition 3.1 (Weak solution). A function $\mathbf{u} : [0, T] \rightarrow H_0^1(\Omega)$ is a weak solution to the problem (1) with the condition (2) if

- (i) $\mathbf{u} \in L^2([0, T]; H_0^1(\Omega))$ and ${}^{RL}D_{0+}^\alpha \mathbf{u} \in L^2([0, T]; H^{-1}(\Omega))$;
- (ii) for all $v \in H_0^1(\Omega)$

$$\langle {}^{RL}D_{0+}^\alpha \mathbf{u}(t), v \rangle_{H^{-1} \times H_0^1} + a(\mathbf{u}(t), v; t) = \langle f(t), v \rangle_{H^{-1} \times H_0^1}$$

for a.e. $t \in (0, T)$, where $\langle \cdot, \cdot \rangle_{H^{-1} \times H_0^1}$ is the duality pairing between $H^{-1}(\Omega)$ and $H_0^1(\Omega)$.

3.1 Galerkin approximation solution

Let $\{e_j\}_{j=1}^{\infty}$ be smooth functions which constitutes an orthonormal basis of $L^2(\Omega)$ and a basis of $H_0^1(\Omega)$ such that

(i) $-\Delta e_k = \lambda_k e_k, k \in \mathbb{N};$

(ii)

$$\int_{\Omega} e_j e_j = \begin{cases} 1, & i = j \\ 0, & i \neq j \end{cases} \quad \text{and} \quad \int_{\Omega} D e_i D e_j = \begin{cases} \lambda_i, & i = j \\ 0, & i \neq j \end{cases}$$

For the existence of these functions, see [E98, p. 335]. Fix $N \in \mathbb{N}$. Let $E_N = \text{span}\{e_1, \dots, e_N\}$ and P_N be the project map from $L^2(\Omega)$ to E_N defined by $P_N u = \sum_{i=1}^N c^i e_i$ with $u \in L^2(\Omega)$ has the form $u = \sum_{i=1}^{\infty} c^i e_i$ in which $c^i = \int_{\Omega} u e_i dx, i \in \mathbb{N}$. In this section, using Galerkin method, we will construct approximation solutions to the problem (1)–(2) in E_N .

Let the function $\mathbf{u}_N : [0, T] \rightarrow E_N$ having the form

$$\mathbf{u}_N(t) = \sum_{i=1}^N c^i(t) e_i, \quad t \in [0, T], \quad (3)$$

here $c^i(\cdot), 1 \leq i \leq N$, is continuous and has the Riemann–Liouville fractional derivative of the order α on $[0, T]$. Assume that

$$({}^{RL}D_{0+}^{\alpha} \mathbf{u}_N, v)_{L^2} + a(\mathbf{u}_N(t), v; t) = \langle f(t), v \rangle_{H^{-1} \times H_0^1} \quad (4)$$

for a.e. $t \in (0, T)$ and

$$\mathbf{u}_N(0) = 0. \quad (5)$$

. We obtain the following result.

Proposition 3.2. For any $N \in \mathbb{N}$, there exists a unique solution to the problem (4)–(5) having the form (3).

Proof. Consider the function $\mathbf{u}_N : [0, T] \rightarrow E_N$ having the form

$$\mathbf{u}_N(t) = \sum_{i=1}^N c^i(t) e_i, \quad t \in [0, T],$$

where $c^i(\cdot), 1 \leq i \leq N$, is continuous and has the Riemann–Liouville fractional derivative of the order α on $[0, T]$. To $\mathbf{u}_N(\cdot)$ is a solution to (4)–(5) then $c^i(\cdot), 1 \leq i \leq N$, have to satisfy the following condition

$$\begin{aligned} \left(\sum_{i=1}^N {}^{RL}D_{0+}^{\alpha} c^i(t) e_i, e_j \right)_{L^2} + a\left(\sum_{i=1}^N c^i(t) e_i, e_j; t \right) &= {}^{RL}D_{0+}^{\alpha} c^j(t) + \sum_{i=1}^N c^i(t) a(e_i, e_j; t) \\ &= \langle f(t), e_j \rangle_{H^{-1} \times H_0^1} = f^j(t) \end{aligned} \quad (6)$$

for $1 \leq j \leq N$ and almost every $t \in (0, T]$. Moreover,

$$c^j(0) = 0, \quad 1 \leq j \leq N. \quad (7)$$

Put $\vec{c}(t) = (c^1(t), \dots, c^N(t))^T$, $A(t) = (a(e_i, e_j; t))_{1 \leq i, j \leq N}$, and

$$\vec{f}(t) = (f^1(t), \dots, f^N(t))^T.$$

Then the system (6)–(7) is rewritten in the form

$$\begin{aligned} {}^{RL}D_{0+}^\alpha \vec{c}(t) + A(t) \vec{c}(t) &= {}^C D_{0+}^\alpha \vec{c}(t) + A(t) \vec{c}(t) = \vec{f}(t), \quad t \in (0, T], \quad (8) \\ \vec{c}(0) &= 0. \quad (9) \end{aligned}$$

Hence, the system (8)–(9) has a solution on $[0, T]$ if and only if the following integral equation has a continuous solution

$$\vec{c}(t) = -\frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} A(\tau) \vec{c}(\tau) d\tau + \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} \vec{f}(\tau) d\tau, \quad t \in [0, T]. \quad (10)$$

On the space $C([0, T]; \mathbb{R}^N)$, we establish an operator as

$$\mathcal{T}_0 \varphi(t) = -\frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} A(\tau) \varphi(\tau) d\tau + \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} \vec{f}(\tau) d\tau, \quad t \in [0, T].$$

For any $\gamma > 0$, define a norm $\|\cdot\|_\gamma$ on $C([0, T]; \mathbb{R}^N)$ by

$$\|\varphi\|_\gamma := \max_{t \in [0, T]} \frac{\|\varphi(t)\|}{\exp(\gamma t)}.$$

It is obvious that $(C([0, T]; \mathbb{R}^N), \|\cdot\|_\gamma)$ is a Banach space. On the other hand, for any $\varphi, \tilde{\varphi} \in C([0, T]; \mathbb{R}^N)$ and $t \in [0, T]$, we have

$$\begin{aligned} \frac{\|\mathcal{T}_0 \varphi(t) - \mathcal{T}_0 \tilde{\varphi}(t)\|}{\exp(\gamma t)} &\leq \frac{\text{ess sup}_{t \in [0, T]} \|A(t)\|}{\Gamma(\alpha)} \int_0^t \frac{\exp(-\gamma(t-\tau))}{(t-\tau)^{1-\alpha}} \frac{\|\varphi(\tau) - \tilde{\varphi}(\tau)\|}{\exp(\gamma \tau)} d\tau \\ &\leq \frac{\text{ess sup}_{t \in [0, T]} \|A(t)\|}{\gamma^\alpha \Gamma(\alpha)} \int_0^{\gamma t} u^{\alpha-1} \exp(-u) du \|\varphi - \tilde{\varphi}\|_\gamma \\ &\leq \frac{\text{ess sup}_{t \in [0, T]} \|A(t)\|}{\gamma^\alpha} \|\varphi - \tilde{\varphi}\|_\gamma. \end{aligned}$$

Hence, for $\gamma > 0$ large enough, the operator \mathcal{T}_0 is contractive in $(C([0, T]; \mathbb{R}^N), \|\cdot\|_\gamma)$ and has a unique fixed point which is also the solution to the system (10). The proof is complete. \square

We now give some estimate of the Galerkin approximation solution.

Proposition 3.3. For any $N \in \mathbb{N}$, there exists a positive constant C , depending on Ω , T and the coefficients of the equation (1), such that

$$\begin{aligned} \|\mathbf{u}_N\|_{L^2([0, T]; L^2(\Omega))}^2 + \|\mathbf{u}_N\|_{L^2([0, T]; H_0^1(\Omega))}^2 + \|{}^{RL}D_{0+}^\alpha \mathbf{u}_N\|_{L^2([0, T]; H^{-1}(\Omega))}^2 \\ \leq C \|f\|_{L^\infty([0, T]; H^{-1}(\Omega))}^2. \end{aligned}$$

Proof. First, using [T18, Theorem 2], we have

$$\begin{aligned}
\langle {}^{RL}D_{0+}^\alpha \mathbf{u}_N(t), \mathbf{u}_N(t) \rangle_{H^{-1} \times H_0^1} &= \langle {}^C D_{0+}^\alpha \mathbf{u}_N(t), \mathbf{u}_N(t) \rangle_{H^{-1} \times H_0^1} \\
&= \left(\sum_{i=1}^N {}^C D_{0+}^\alpha c^i(t) e_i, \sum_{i=1}^N c^i(t) e_i \right)_{L^2} \\
&= \sum_{i=1}^N c^i(t) {}^C D_{0+}^\alpha c^i(t) \\
&\geq \frac{1}{2} \sum_{i=1}^N {}^C D_{0+}^\alpha (c^i(t))^2 \\
&= \frac{1}{2} {}^C D_{0+}^\alpha \sum_{i=1}^N (c^i(t))^2 \\
&= \frac{1}{2} {}^C D_{0+}^\alpha \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2.
\end{aligned}$$

On the other hand, by [E98, Theorem 3, p. 300], there exist $\beta > 0$ and $\nu \geq 0$ such that

$$\beta \|\mathbf{u}_N(t)\|_{H_0^1}^2 \leq a(\mathbf{u}_N(t), \mathbf{u}_N(t); t) + \nu \|\mathbf{u}_N(t)\|_{L^2}^2 \text{ for almost every } t \in (0, T).$$

Moreover, from the assumption of f and the Cauchy inequality

$$\begin{aligned}
\langle f(t), \mathbf{u}_N(t) \rangle_{H^{-1} \times H_0^1} &\leq \|f(t)\|_{H^{-1}(\Omega)} \|\mathbf{u}_N(t)\|_{H_0^1(\Omega)} \\
&\leq \frac{1}{4\beta} \|f(t)\|_{H^{-1}(\Omega)}^2 + \beta \|\mathbf{u}_N(t)\|_{H_0^1(\Omega)}^2.
\end{aligned}$$

Thus, for almost every $t \in (0, T)$,

$${}^C D_{0+}^\alpha \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2 \leq 2\nu \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2 + \frac{1}{2\beta} \|f(t)\|_{H^{-1}(\Omega)}^2.$$

Put $v(t) := \|\mathbf{u}_N(t)\|_{L^2}^2$, $h(t) := \frac{1}{2\beta} \|f(t)\|_{H^{-1}}^2$ and use the comparison principle for solutions to fractional differential equation and the variation of constants formula for solutions to the equations (see [CT17, Lemma 3.1]), we obtain the estimate

$$v(t) \leq \int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}(2\nu(t-\tau)^\alpha) h(\tau) d\tau \text{ for a.a. } t \in [0, T].$$

This implies that

$$\|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2 \leq \frac{T^\alpha E_{\alpha,\alpha+1}(2\nu T^\alpha) \|f\|_{L^\infty(0,T;H^{-1}(\Omega))}^2}{2\beta} \text{ for a.a. } t \in [0, T]. \quad (11)$$

Next, by the similar arguments as above, we see that

$${}^C D_{0+}^\alpha \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2 + \beta \|\mathbf{u}_N(t)\|_{H_0^1(\Omega)}^2 \leq \frac{1}{\beta} \|f(t)\|_{H^{-1}(\Omega)}^2 + 2\nu \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2.$$

Furthermore,

$${}^C D_{0+}^\alpha \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2 = {}^{RL} D_{0+}^\alpha \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2 = \frac{d}{dt} I_{0+}^{1-\alpha} (\|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2),$$

$$\begin{aligned} & I_{0+}^{1-\alpha} (\|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2) \Big|_0^T + \beta \|\mathbf{u}_N\|_{L^2(0,T;H_0^1(\Omega))}^2 \\ & \leq \frac{T \|f\|_{L^\infty(0,T;H^{-1}(\Omega))}^2}{\beta} + 2\nu \int_0^T \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2 dt, \end{aligned}$$

Thus,

$$\beta \|\mathbf{u}_N\|_{L^2([0,T];H_0^1(\Omega))}^2 \leq \frac{T \|f\|_{L^\infty([0,T];H^{-1}(\Omega))}^2}{\beta} + 2\nu \int_0^T \|\mathbf{u}_N(t)\|_{L^2(\Omega)}^2 dt.$$

This combines with (11) implies that there is a constant $C_1 > 0$ such that

$$\|\mathbf{u}_N\|_{L^2([0,T];H_0^1(\Omega))}^2 \leq C_1 \|f\|_{L^\infty([0,T];H^{-1}(\Omega))}^2. \quad (12)$$

Finally, fix any $v \in H_0^1(\Omega)$ with $\|v\|_{H_0^1} \leq 1$, and write $v = v_0 + v_1$, where $v_1 \in \text{span}\{e_1, \dots, e_N\}$ and $(e_i, v_0) = 0$, $1 \leq i \leq N$. Using the estimate concerning the bilinear operator $a(\cdot, \cdot)$ and the facts that $({}^{RL} D_{0+}^\alpha u_N(t), v_1)_{L^2} = \langle {}^{RL} D_{0+}^\alpha u_N(t), v \rangle_{H^{-1} \times H_0^1}$, $({}^{RL} D_{0+}^\alpha u_N(t), v_1)_{L^2} + a(u_N(t), v_1; t) = \langle f(t), v_1 \rangle_{H^{-1} \times H_0^1}$ and $\|v_1\|_{H_0^1(\Omega)} \leq \|v\|_{H_0^1(\Omega)} \leq 1$, we obtain

$$\begin{aligned} |\langle {}^{RL} D_{0+}^\alpha \mathbf{u}_N(t), v \rangle_{H^{-1} \times H_0^1}| & \leq |a(\mathbf{u}_N(t), v_1; t)| + |\langle f(t), v_1 \rangle_{H^{-1} \times H_0^1}| \\ & \leq C_2 \|\mathbf{u}_N(t)\|_{H_0^1(\Omega)} \|v\|_{H_0^1(\Omega)} + \|f(t)\|_{H^{-1}(\Omega)} \|v\|_{H_0^1(\Omega)}. \end{aligned}$$

Hence,

$$\|{}^{RL} D_{0+}^\alpha \mathbf{u}_N(t)\|_{H^{-1}(\Omega)} \leq C_2 \|\mathbf{u}_N(t)\|_{H_0^1(\Omega)} + \|f(t)\|_{H^{-1}(\Omega)}$$

which together with (13) and the estimate (12) implies

$$\|{}^{RL} D_{0+}^\alpha \mathbf{u}_N\|_{L^2([0,T];H^{-1}(\Omega))}^2 \leq C_3 \|f\|_{L^\infty([0,T];H^{-1}(\Omega))}^2. \quad (13)$$

From (11), (12) and (13), the proof is complete. \square

3.2 Existence and uniqueness of weak solutions

We now in a position to state the main result of the paper.

Theorem 3.4. Consider the problem (1)–(2). Suppose that assumptions (a1)–(a4) hold. Then, this problem has a unique weak solution.

Proof. First, we prove the system (1)–(2) has at least one weak solution. From (12) and (13), by the Banach–Alaoglu theorem, there exist a sequence $\{n_k\}_{k=1}^\infty$ such that

$$\mathbf{u}_{n_k} \rightharpoonup u \text{ in } L^2_0([0, T]; H^1_0(\Omega)), \quad (14)$$

$${}^{RL}D_{0+}^\alpha \mathbf{u}_{n_k} \rightharpoonup v \text{ in } L^2([0, T]; H^{-1}(\Omega)). \quad (15)$$

Let $\varphi \in C_c^\infty((0, T); \mathbb{R})$ and $\psi \in H^1_0(\Omega)$ be arbitrary. Then,

$$\begin{aligned} \int_0^T \langle v(t), \phi(t)\psi \rangle_{H^{-1} \times H^1_0} dt &= \lim_{k \rightarrow \infty} \int_0^T \langle {}^{RL}D_{0+}^\alpha \mathbf{u}_{n_k}(t), \phi(t)\psi \rangle_{H^{-1} \times H^1_0} dt \\ &= \lim_{k \rightarrow \infty} \int_0^T \phi(t) \langle {}^{RL}D_{0+}^\alpha \mathbf{u}_{n_k}(t), \psi \rangle_{H^{-1} \times H^1_0} dt \\ &= \lim_{k \rightarrow \infty} \int_0^T {}^{RL}D_{T-}^\alpha \phi(t) \langle \mathbf{u}_{n_k}(t), \psi \rangle_{H^{-1} \times H^1_0} dt \\ &= \int_0^T {}^{RL}D_{T-}^\alpha \phi(t) \langle \mathbf{u}(t), \psi \rangle_{H^{-1} \times H^1_0} dt \\ &= \int_0^T \langle {}^{RL}D_{0+}^\alpha \mathbf{u}(t), \varphi(t)\psi \rangle_{H^{-1} \times H^1_0} dt \\ &= \int_0^T \varphi(t) \langle {}^{RL}D_{0+}^\alpha \mathbf{u}(t), \psi \rangle_{H^{-1} \times H^1_0} dt, \end{aligned}$$

which implies

$$v(t) = {}^{RL}D_{0+}^\alpha \mathbf{u}(t). \quad (16)$$

Fix $N, M \in \mathbb{N}$ and $N > M$. For any $\varphi \in C_0^\infty([0, T]; \mathbb{R})$ and $w \in E_M$, we see that

$$\begin{aligned} \int_0^T \langle {}^{RL}D_{0+}^\alpha \mathbf{u}_N(t), \varphi(t)w \rangle_{H^{-1} \times H^1_0} dt &= \int_0^T \varphi(t) \langle {}^{RL}D_{0+}^\alpha \mathbf{u}_N(t), w \rangle_{H^{-1} \times H^1_0} dt \\ &\rightarrow \int_0^T \varphi(t) \langle {}^{RL}D_{0+}^\alpha \mathbf{u}(t), w \rangle_{H^{-1} \times H^1_0} dt, \end{aligned}$$

$$\begin{aligned} \int_0^T a(\mathbf{u}_N(t), \varphi(t)w; t) dt &= \int_0^T \varphi(t) a(\mathbf{u}_N(t), w; t) dt \\ &\rightarrow \int_0^T \varphi(t) a(\mathbf{u}(t), w; t) dt, \end{aligned}$$

and

$$\int_0^T \langle f(t), \varphi(t)w \rangle_{H^{-1} \times H^1_0} dt = \int_0^T \varphi(t) \langle f(t), w \rangle_{H^{-1} \times H^1_0} dt.$$

Thus,

$$\langle {}^{RL}D_{0+}^\alpha \mathbf{u}(t), w \rangle_{H^{-1} \times H^1_0} + a(\mathbf{u}(t), w; t) = \langle f(t), w \rangle_{H^{-1} \times H^1_0}. \quad (17)$$

By combining (14), (15), (16) and (17), there exists a unique weak solution to the problem (1)–(2).

Next, we give a proof for the uniqueness of weak solutions to (1)–(2). Suppose that u_1, u_2 are two weak solution to this system. Define $u := u_1 - u_2$. Then, u satisfies

$$\langle (k * u)'(t), v \rangle_{H^{-1} \times H_0^1} + a(u(t), v; t) = 0, \quad v \in H_0^1(\Omega), \quad a.a. \ t \in (0, T).$$

Let $v = u(t)$, then for almost every $t \in (0, T)$

$$\begin{aligned} \langle {}^{RL}D_{0+}^\alpha u(t), u(t) \rangle_{H^{-1} \times H_0^1} + a(u(t), u(t); t) &= \langle (k * u)'(t), u(t) \rangle_{H^{-1} \times H_0^1} \\ &+ a(u(t), u(t); t) = 0, \end{aligned} \quad (18)$$

where $k(t) = \frac{1}{\Gamma(1-\alpha)t^\alpha}$ for $t > 0$. Motivated by Rico Zacher [VZ08, pp. 291–292], we will approximate the operator $\frac{d}{dt}(k * u)$ by the sequences $\{\frac{d}{dt}(k_n * u)\}_n$, where $k_n(t) := ns(t) = nE_\alpha(-nt^\alpha)$ (note that by using Laplace transform, we see that $s(\cdot)$ is the unique solution to the equation $s(t) + n(l * s)(t) = 1$, $t > 0$, where $l(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$, $t > 0$, see [KTT20, p. 3]). We rewrite the equation (18) as below

$$\langle (k_n * u)'(t), u(t) \rangle_{H^{-1} \times H_0^1} + a(u(t), u(t); t) = h_n(t), \quad a.a. \ t \in (0, T) \quad (19)$$

with

$$h_n(t) := \langle (k_n * u)'(t) - (k * u)'(t), u(t) \rangle_{H^{-1} \times H_0^1}, \quad a.a. \ t \in (0, T).$$

By virtue of [Z09, Lemma 2.1],

$$\frac{1}{2} \frac{d}{dt} (k_n * \|u(\cdot)\|_{L^2(\Omega)}^2)(t) \leq \left(\frac{d}{dt} (k_n * u)(t), u(t) \right)_{L^2(\Omega)}, \quad a.a. \ t \in (0, T).$$

On the other hand, there exists $d \geq 0$ such that

$$a(u(t), u(t); t) \geq -\nu \|u(t)\|_{L^2(\Omega)}^2.$$

Hence, from (19), we have

$$\frac{d}{dt} (k_n * \|u(\cdot)\|_{L^2(\Omega)}^2)(t) \leq 2\nu \|u(t)\|_{L^2(\Omega)}^2 + 2h_n(t), \quad a.a. \ t \in (0, T).$$

This together with the positivity of l implies that

$$l * \frac{d}{dt} (k_n * \|u(\cdot)\|_{L^2(\Omega)}^2) \leq 2\nu (l * \|u(\cdot)\|_{L^2(\Omega)}^2)(t) + 2l * h_n(t), \quad a.a. \ t \in (0, T).$$

We will show that

$$\lim_{n \rightarrow \infty} h_n = 0 \text{ in } L^1([0, T]; \mathbb{R}). \quad (20)$$

From the facts that $u \in L^2([0, T]; H^{-1}(\Omega))$, ${}^{RL}D_{0+}^\alpha u = \frac{d}{dt}[k*u] \in L^2([0, T]; H^{-1}(\Omega))$, [VZ08, Proposition 2.1, p. 293] and [VZ08, Example 2.1, p. 294], we have

$$k * \|u(\cdot)\|_{H^{-1}(\Omega)}^2 \in W_1^1([0, T]; \mathbb{R}).$$

This implies that

$$u \in D(B_2), \|u(\cdot)\|_{H^{-1}(\Omega)}^2 \in D(B_1),$$

where

$$B_1(u) = \frac{d}{dt}k*u, D(B_1) = \{u \in L^1(0, T) : k*u \in W_1^1([0, T]; \mathbb{R})\},$$

and

$$B_2(u) = \frac{d}{dt}k*u, D(B_2) = \{u \in L^2([0, T]; H^{-1}(\Omega)) : k*u \in W_2^1([0, T]; H^{-1}(\Omega))\}.$$

Hence, by using [VZ08, Estimate (18), p. 292], we obtain

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_0^T \left\| \frac{d}{dt}[(k - k_n) * u](t) \right\|_{H^{-1}(\Omega)}^2 dt \\ &= \lim_{n \rightarrow \infty} \left\| \frac{d}{dt}[(k - k_n) * u](\cdot) \right\|_{L^2([0, T]; H^{-1}(\Omega))}^2 = 0. \end{aligned} \quad (21)$$

By (21) and the Holder inequality, the following estimates hold

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_0^T |h_n(t)| dt &= \lim_{n \rightarrow \infty} \int_0^T | \langle (k_n * u)'(t) - (k * u)'(t), u(t) \rangle_{H^{-1} \times H_0^1} | dt \\ &\leq \lim_{n \rightarrow \infty} \int_0^T \left\| \frac{d}{dt}[(k - k_n) * u](t) \right\|_{H^{-1}(\Omega)} \|u(t)\|_{H_0^1(\Omega)} dt \\ &\leq \lim_{n \rightarrow \infty} \left\| \frac{d}{dt}[(k - k_n) * u](\cdot) \right\|_{L^2([0, T]; H^{-1}(\Omega))} \|u\|_{L^2([0, T]; H_0^1(\Omega))} \\ &= 0, \end{aligned}$$

which shows that $\lim_{n \rightarrow \infty} h_n = 0$ in $L^1([0, T])$. Notice that

$$\begin{aligned} (l * k)(t) &= \frac{1}{\Gamma(\alpha)\Gamma(1-\alpha)} \int_0^t s^{\alpha-1}(t-s)^{-\alpha} ds \\ &= \frac{1}{\Gamma(\alpha)\Gamma(1-\alpha)} \int_0^1 u^{-\alpha}(1-u)^{\alpha-1} du \\ &= \frac{1}{\Gamma(\alpha)\Gamma(1-\alpha)} B(\alpha, 1-\alpha) \\ &= 1, \quad \forall t > 0, \end{aligned}$$

$$l * \frac{d}{dt}(k_n * \|u(\cdot)\|_{L^2(\Omega)}^2) = \frac{d}{dt}(k_n * l * \|u(\cdot)\|_{L^2(\Omega)}^2) \rightarrow \frac{d}{dt}(k * l * \|u(\cdot)\|_{L^2(\Omega)}^2) = \|u(\cdot)\|_{L^2(\Omega)}^2$$

in $L^1([0, T]; \mathbb{R})$ as $n \rightarrow \infty$ (see [VZ08, Estimate (19), p. 292]) and $l * h_n \rightarrow 0$ in $L^1([0, T]; \mathbb{R})$ as $n \rightarrow \infty$, we obtain

$$\|u(t)\|_{L^2(\Omega)}^2 \leq 2\nu(l * \|u(\cdot)\|_{L^2(\Omega)}^2)(t), \text{ a.a. } t \in (0, T). \quad (22)$$

From (22), using a Gronwall type inequality as in [H81, Lemma 7.1.1, p. 188], then $\|u(t)\|_{L^2(\Omega)}^2 = 0$ a.e. in $(0, T)$, that is, $u_1 = u_2$. The proof is complete. \square

Remark 3.5. The key point in the proof of the uniqueness of weak solution to the problem (1)–(2) is to show that $h_n \rightarrow 0$ in $L^1([0, T])$ as $n \rightarrow \infty$. The approach proposed in [Z09] can not apply directly to this situation because the operator $B = \frac{d}{dt}(k * u)$ with domain $D(B) = \{u \in L^2([0, T]; H_0^1(\Omega)) : \frac{d}{dt}k * u \in L^2([0, T]; H^{-1}(\Omega))\}$ is not m-accretive.

Acknowledgement

A part of this paper was completed while the author was a postdoc at the Vietnam Institute for Advanced Study in Mathematics (VIASM). He would like to thank VIASM for support and hospitality.

References

- [B20] K.V. Bockstal, Existence and uniqueness of a weak solution to a non-autonomous time-fractional diffusion equation (of distributed order). *Applied Mathematics Letters*, **109** (2020), 106540.
- [C99] M. Caputo, Diffusion of fluids in porous media with memory. *Geothermics*, **28** (1999), pp. 113–130.
- [CT17] N.D. Cong and H.T. Tuan, Generation of nonlocal fractional dynamical systems by fractional differential equations. *Journal of Integral Equations and Applications*, **29** (2017), no. 4, pp. 585–608.
- [EK04] S.D. Eidelman, A.N. Kochubei, Cauchy problem for fractional diffusion equations. *Journal of Differential Equations*, **199** (2004), no. 2, pp. 211–255.
- [E98] L.C. Evans, *Partial Differential Equations*. Graduate Series in Mathematics, **19**. American Mathematics Society, 1998.
- [H81] D. Henry, *Geometric Theory of Semi-linear Parabolic Equations*. Lecture Notes in Mathematics, **840**. Springer, 1981.
- [GLY15] R. Gorenflo, Y. Luchko, and M. Yamamoto, Time-fractional diffusion equation in the fractional Sobolev spaces. *Fractional Calculus and Applied Analysis*, **18** (2015), no. 3, pp. 799–820.

- [KTT20] T.D. Ke, N.N. Thang, L.T.P. Thuy, Regularity and stability analysis from a class of semilinear nonlocal differential equations in Hilbert spaces. *Journal of Mathematical Analysis and Applications*, **483** (2020), 123655.
- [LY37] E. Love and L. Young, On fractional integration by parts. *Proceedings of The London Mathematical Society*, **44** (1937), 1–35.
- [MK2000] R. Metzler, J. Klafter, Boundary value problems for fractional diffusion equations. *Physica A: Statistical Mechanics and its Applications*, **278** (2000), no. 1–2, pp. 107–125.
- [N86] R.R. Nigmatullin, The realization of the generalized transfer equation in a medium with fractal geometry. *Physica Status Solidi (b)*, **133** (1986), no. 1, pp. 425–430.
- [P93] J. Pruss, *Evolutionary Integral Equations and Applications*. Monograph in Mathematics **87**, Birkhauser, Basel, 1993.
- [SKM93] S.G. Samko, A.A. Kilbas, and O.I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*. Gordon and Breach Science Publishers, 1993.
- [T18] H.T. Tuan, H. Trinh, Stability of fractional-order nonlinear systems by Lyapunov direct method. *IET Control Theory and Applications*, **12** (2018), pp. 2417–2422.
- [RA94] H.E. Roman, P.A. Alemany, Continuous-time random walks and the fractional diffusion equation. *Journal of Physics A: Mathematical and General*, **27** (1994), no. 10, pp. 3407–3410.
- [SY11] K. Sakamoto, M. Yamamoto, Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems. *Journal of Mathematical Analysis and Applications*, **382** (2011), no. 1, pp. 426–447.
- [VZ08] V. Vergara and R. Zacher, Lyapunov functions and convergence to steady state for differential equations of fractional order. *Math.Z.*, **259** (2008), pp. 287–309.
- [Z09] R. Zacher, Weak solutions of abstract evolutionary integro-differential equations in Hilbert spaces. *Funkcialj Ekvacioj*, **52** (2009), pp. 1–18.